

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Expression and functionality of TRPV1 receptor in human MCF-7 and canine CF.41 cells.

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/137174> since 2021-11-17T22:38:03Z

Published version:

DOI:10.1111/vco.12028

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Expression and functionality of TRPV1 receptor in human MCF-7 and canine CF.41 cells

C. Vercelli, R. Barbero, B. Cuniberti, R. Odore and G. Re

Department of Veterinary Sciences, Division of Veterinary Pharmacology and Toxicology, University of Turin, Grugliasco, Turin, Italy

Correspondence address:

C. Vercelli

Department of Veterinary Sciences

Division of Veterinary Pharmacology and Toxicology University of Turin

Via Leonardo da Vinci 44 I-10095 Grugliasco Turin, Italy

e-mail: cristina.vercelli@unito.it

Abstract

As canine mammary tumours (CMT) and human breast cancer share clinical and prognostic features, the former have been proposed as a model to study carcinogenesis and improved therapeutic treatment in human breast cancer. In recent years, it has been shown that transient receptor potential vanilloid 1 (TRPV1) is expressed in different neoplastic tissues and its activation has been associated with regulation of cancer growth and progression. The aim of the present research was to demonstrate the presence of TRPV1 in human and canine mammary cancer cells, MCF-7 and CF.41, respectively, and to study the role of TRPV1 in regulating cell proliferation. The images obtained by Western blot showed a signal at 100 kDa corresponding to the molecular weight of TRPV1 receptor. All tested TRPV1 agonists and antagonists caused a significant decrease ($P < 0.05$) of cell growth rate in MCF-7 cells. By contrast, in CF.41 cells capsaicin and capsazepine induced a significant increase

($P < 0.05$) in cell proliferation, whereas resiniferatoxin (RTX) and 5-iodo-resiniferatoxin (5-I-RTX) had

no influence on CF.41 cell proliferation. Further studies are needed to elucidate the underlying molecular mechanism responsible for the different effects evoked by TRPV1 activation in MCF-7 and CF.41 cells.

Keywords

CF.41, MCF-7, proliferation assays, TRPV1, Western blotting

Introduction

Recent epidemiological data suggest that in 2008 almost 1.4 million women worldwide were diagnosed with breast cancer and approximately 459 000 deaths were recorded. The 5-year relative survival rate ranges from 12 to 90% depending on diagnostic and therapeutic approach.¹ Similarly to humans, mammary glands are the most frequent site of neoplasia in female dogs and in this species mammary tumours represent one of the leading causes of death due to development of metastases in regional lymph nodes and lung.² Clinical and molecular similarities between canine mammary tumours (CMT) and human breast cancer have been described in recent years.³ Interestingly, beside the fact that the etiopathogenesis of CMT is still unclear, by comparing breast cancer frequencies of dog and pet ownership a significant positive correlation was found.⁴ Indeed, the use of dogs spontaneously affected by cancer in trials testing novel anticancer therapies for humans, is growing up.^{5,6} As in humans, age, obesity and diet seem to increase the risk of CMT.⁷ Moreover, in both species prognosis depends on tumour size, histology type, proliferation rate, lymph node status, hormonal receptor status, presence of metastases and gene expression profile.⁸

The transient receptor potential vanilloid 1 (TRPV1) is a member of the transient receptor potential (TRP) family known to be expressed in several neural and non-neuronal tissues of different animal species.⁹ TRPV1 is a non-selective cation channel that can be activated either by physical, chemical and pro-inflammatory stimuli such as noxious heat (>43 °C), low pH (<6) or by different lipophilic compounds including capsaicin and phospholipid derivatives.¹⁰ The activation of TRPV1 receptors causes calcium influx and the corresponding release of tachykinin neuropeptides which, in turn, trigger a tissue response known as 'neurogenic' inflammation.¹¹ In recent years, TRP channels have been associated with cancer growth and progression.¹² Their expression has been demonstrated in different neoplastic tissues such as squamous cell carcinoma of the human tongue,¹³ human prostate carcinoma,¹⁴ colon cancer,¹⁵ transitional carcinoma of human bladder¹⁶ and human breast cancer.¹⁷ Santoni et al.¹⁶ found that TRPV1 expression decreases during the development of transitional cell carcinoma of human bladder and outlined its key role in the regulation of tumour cell proliferation and even as possible therapeutic target.¹⁶ Mammalian TRPV1 receptors were cloned and characterized in humans,¹⁸ rats,¹⁹ guinea pigs,²⁰ rabbits,²¹ mice²² and dogs.²³ The various TRPV1-homologues share a conserved structure composed by six transmembrane spanning domains (S1 – S6), intracellular N and C terminals and a pore-forming hydrophobic stretch between the fifth and the sixth domains. TRPV1 receptors are activated by vanilloids-like capsaicin. However, TRPV1 proteins do not always exhibit the same pharmacological profile. While in rat 12-phenyl-acetate 13-acetate 20-homovanillate (PPAHV) acts as a potent agonist for TRPV1, in humans and guinea pig TRPV1s are not stimulated by PPAHV.^{24,25}

As mentioned previously, the canine model is currently used in comparative oncology as spontaneous tumours in dogs share a wide variety of epidemiological, biological and clinical features with human cancers, which make this animal model attractive in oncology research.²⁶ In past decades, dog was studied as a model for prostate tumour,²⁷ osteosarcoma,²⁸ lymphoma^{29,30} and mammary tumour.^{26,31} Considering that TRPV1 is expressed both in human and canine species and it is expressed in human breast cancer, the aim of the present research was to study the expression of TRPV1 receptor in cells derived from human breast cancer and canine mammary cancer. Moreover, the pharmacological characteristics of different specific TRPV1 agonists and antagonists were studied through proliferation assays in order to evaluate the capacity of TRPV1 ligands to modulate the growth of tumour cells.

Materials and methods

Cell culture

MCF-7 cells (cells derived from pleural effusion after pleural metastatization of human breast adenocarcinoma, ATCC cell culture, LGC Standards, Milan, Italy) and CF.41 cells (cells derived from canine mammary adenocarcinoma, ATCC cell culture, LGC Standards, Milan, Italy) were cultured in 75 cm³ flasks containing Dulbecco's Modified Eagles Medium (DMEM, Lonza, Basel, Switzerland) supplemented with 20% Fetal Bovine Serum (FBS, Sigma Aldrich, Milan, Italy), 2% penicillin–streptomycin–amphotericin B solution (Sigma Aldrich) and 2% L-glutamine (Sigma Aldrich) at 37 °C with 5% CO₂, 95% air and complete humidity. When reaching a confluence of 80%, cells were detached using 0.1% Trypsin/EDTA solution (Sigma Aldrich), centrifuged and counted using Trypan Blue solution (Sigma Aldrich) with a Burkert chamber. Cells were then resuspended at a concentration of 5 × 10⁵ cells mL⁻¹ and seeded at the same conditions mentioned before.

Western blotting assay

Thawed cells (concentration 180 × 10⁶ cells approximately), re-suspended in binding buffer (KCl 5 mM, NaCl 5.8 mM, CaCl₂ 0.75 mM, MgCl₂ 2 mM, sucrose 320 mM and Hepes 10 mM, pH 7.4, Sigma Aldrich), were disrupted using a sonicator (XL 2020, Misonix, Farmingdale, NY, USA) and centrifuged at 3000 × g for 10 min at 4 °C. The resulting supernatants were ultra centrifuged at 105 000 × g for 45 min at 4 °C to separate

cytosolic fraction from cell membrane fraction. The final pellets were suspended in the same buffer and the protein concentration was measured according to the method described by Lowry et al.³² Samples were loaded at 25, 50 and 100 μ g mL⁻¹ of protein per lane and subjected to sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) procedure using 8% Acrylamide (Biosolve, Valkenswaard, The Netherlands) gel. A proteic lysate of urinary rat bladder was used as positive control. Samples were then transferred to nitrocellulose membranes (Hybond ECL, Amersham, GE Healthcare, Milan, Italy). After being washed three times, samples were probed with the primary anti-TRPV1 antibody (goat polyclonal antibody, Santa Cruz Biotechnology, Heidelberg, Germany) (dilution 1:100) and incubated overnight at 5 °C on a shaking plate. At the end of the incubation period samples were washed again for four times and probed with the secondary antibody (donkey anti-goat IgG-HRP, Santa Cruz Biotechnology) (dilution 1:15 000) for 1 h. Secondary antibody was removed by four washes and immunoreactive bands were visualized by enhanced chemiluminescence (ECL-Plus System, Amersham, GE Healthcare). To obtain an endogenous control, membranes were washed and re-probed with β -actin antibody (dilution 1:100, Santa Cruz Biotechnology) for 1 h. Also, in this case the immunoreactive bands of β -actin were visualized by ECL (ECL-Plus System, Amersham, GE Healthcare).

Proliferation assays

3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT, Sigma Aldrich) reduction assay was performed for both cell lines. This method is one of the most frequently used for quantitating cell viability and it measures the conversion of MTT into purple-coloured formazan that is induced by redox activity of living cells: a decrease of cellular redox activity could be index of cell damage.³³

Cultured cells were detached using 0.1% Trypsin/EDTA solution (Sigma Aldrich) and diluted to a final concentration of 5×10^3 cells 100 μ L⁻¹. The cell suspension was transferred into 96-well plates with DMEM (Lonza, Basel, Switzerland), 10% FBS (Sigma Aldrich), 2% penicillin-streptomycin-amphotericin B solution (Sigma Aldrich) and 2% L-glutamine (Sigma Aldrich). Each drug was tested in two different experimental sessions and three plates were prepared per each experimental session (n = 6). To evaluate the background adsorption, the first row of every plate remained cell-free as blank control.

In the second row, cells were seeded in absence of any drug as internal control. The remaining part of the plates was used to test TRPV1 ligands. After an overnight incubation at 37 °C and 5% CO₂, the medium used for the seeding was replaced by a medium prepared with DMEM, 5% FBS, 2% penicillin-streptomycin-amphotericin B solution, 2% L-glutamine with the addition of increasing concentrations of the following TRPV1 ligands: Capsaicin and capsazepine (partial agonist and antagonist, respectively, Sigma Aldrich) (10⁻¹⁰–10⁻⁴M), resiniferatoxin (RTX) and 5-iodo-resiniferatoxin (5-I-RTX) (pure agonist and antagonist, respectively, Sigma Aldrich) (10⁻¹⁰–10⁻⁶M). Plates were incubated at 37 °C at 5% CO₂ for 24, 48 and 72h. At each experimental time point, 20 μ L of MTT (Sigma Aldrich) solution (4 mg mL⁻¹) were added to each well, except to the free-cell blank wells, and plates were incubated at 37 °C and 5% CO₂ for 4h. At the end of the incubation period, the medium containing the MTT solution was discarded and 100 μ L of dimethyl sulfoxide (DMSO, Sigma Aldrich) were added in order to dissolve the formazan crystals. The plates were further incubated for 10 min at room temperature and then read using a plate reader (Powerwave X, Bio-Tek, Winooski, VT, USA) at a wavelength of 570 nm.

The results obtained by the proliferation assays were statistically analysed using a computer programme (GraphPad Prism Software, San Diego, CA, USA), applying the Kruskal-Wallis test and a Dunn's post-test (P < 0.05).

Results

Western blotting

Western blotting analysis indicated that the TRPV1 protein is expressed in MCF-7 cells as shown by the high density signal corresponding to the molecular weight of 100 kDa, in accordance with the mass of human TRPV1 (Fig. 1A).³⁴ The signal corresponding to 100 kDa was also recognizable in the Western blotting analysis of CF.41 cells (Fig. 1B). The same signal was present in positive control (lysate of rat urinary bladder). Fig. 1C shows the signal corresponding to β -actin, used as endogenous control.

Proliferation assays

MCF-7 proliferation assay data are summarized in Figs 2– 5. All tested agonists and antagonists induced a significant decrease ($P < 0.05$) of growth rate in treated cells compared with controls. RTX showed a characteristic time-dependent behaviour (Fig. 4). As far as CF.41 cells are concerned results are presented in Figs 6– 9. The partial agonist (capsaicin) and the partial antagonist (capsazepine) caused a significant increase ($P < 0.05$) of CF.41 cell proliferation. By contrast, RTX and 5-I-RTX did not induce any significant effect on cell growth rate.

Discussion

To our knowledge, this is the first study demonstrating the expression of TRPV1 in CF.41 cells, while TRPV1 in MCF-7 cells had already been identified by Barbero et al.³⁵ in 2006. Increased expression of TRPV1 is associated with a better prognosis of patients with hepatocellular carcinoma, whereas expression of TRPV1 is decreased in the urothelium of patients with advanced stage transitional cell carcinoma, suggesting that TRPV1 might function as a tumour suppressor and negatively control tumour progression.³⁶ Different TRP channels are also expressed in human breast cancer cells and cell lines such as MCF-7 and their overexpression compared to normal gland can be considered a suitable prognostic marker.³⁷ It has been suggested that the receptor-mediated anti-tumour activity is associated with induction of apoptosis following activation of TRPV1 and changes in cell calcium influx.³⁸ Calcium is an important intracellular messenger that regulates many physiological functions such as cellular differentiation, proliferation and apoptosis. Abnormal functioning of calcium signalling pathways may occur during cancerogenesis. Therefore, calcium channels with increased expression and/or activity in cancer could potentially serve as anti-cancer therapeutic targets. Recent investigations have suggested that besides proapoptotic action, the activation of TRPV1 leads to anti-angiogenic, anti-migrative, anti-adhesive and anti-metastatic effects.³⁹ As TRPV1 is implicated in cell proliferation and pain modulation, a number of studies have been performed to describe the binding characteristics and clinical efficacy of selective TRPV1 agonists and antagonists.⁴⁰ Capsaicin and RTX are selective TRPV1 agonists and cause TRPV1 desensitization after activation while capsazepine and 5-iodoresiniferatoxin are selective TRPV1 antagonists inducing a functional block of TRPV1 receptor.⁴¹ In our study, all tested agonists and antagonists caused a significant decrease ($P < 0.05$) in MCF-7 cell proliferation. By contrast, capsaicin and capsazepine induced a significant increase ($P < 0.05$) in CF.41 cell proliferation. These findings might account for species-specific differences in TRPV1 pharmacological properties. In rodents, TRPV1 is activated by capsaicin and residues Arg114 and Glu761 in the intracellular N- and C-termini have been recognized as agonist binding sites.⁴² Unlike the murine one, the avian form of TRPV1 is not activated by capsaicin and the agonist binding sites are represented by residues Tyr511 and Thr550 in the third and fifth transmembrane domains.²¹ Moreover, species-dependent behaviour has been observed also for antagonists. According to Premkumar and Sikand,⁴³ the TRPV1 competitive antagonist capsazepine inhibits the acid-induced activation of human TRPV1, but exhibits no effects on acid-induced activation of rat TRPV1. The compound can inhibit heat-induced TRPV1 activation both in humans and rat. Besides this, it has been demonstrated that TRPV1 mutations can lead to changes in TRPV1 responses to agonist stimulation. Mutations at the sixth TRPV1 transmembrane domain can cause, for example, the reduction of ^3H resiniferatoxin-binding affinity and mutations on pore domain significantly decrease the sensitivity to capsaicin.⁴⁴ A further hypothesis to explain the different effects induced by TRPV1 agonists and antagonists on MCF-7 and CF.41

proliferation might be the possible interaction of capsaicin and capsazepine with other TRP or non-TRP receptors. Hwang et al.⁴¹ found that capsaicin might act as co-carcinogen on 12-O-tetradecanoylphorbol-13-acetate (TPA)- induced skin carcinogenesis through the epidermal growth factor receptor. As the MTT proliferation assay does not provide information concerning the molecular mechanisms involved in TRPV1 responses, further studies are needed to understand differences in human and canine breast cancer cell proliferation upon TRPV1 agonist and antagonist stimulation. The results of proliferation assays performed on MCF-7 confirm the potential of TRPV1 agonists as anti-cancer drugs. Interestingly, capsaicin was shown to inhibit human cancer hormone-resistant cells and to provide chemopreventive effects.⁴⁵ Moreover, it was demonstrated that capsaicin is able to mediate cell death of T24 (human bladder cancer) and A172 (human glioma) cell lines.^{46,47} Even if the results of proliferation assays on MCF-7 cells seem to suggest that also TRPV1 antagonists could be used in cancer treatment, their use should be carefully considered: the systemic administration of TRPV1 antagonists could lead to the change of body temperature, gastrointestinal and cardiovascular functions.⁴³ Furthermore, the blockade of TRPV1 has been shown to increase its own expression, thereby raising the possibility of rebound effects.⁴³ Therefore, TRPV1-mediated effects on human and canine breast cancer cells should be encouraged.

Acknowledgement

Supported by grants of Regione Piemonte ('Ricerca Sanitaria Finalizzata Regione Piemonte 2009')

References

1. Youlten DR, Cramb SM, Dunn NA, Muller JM, Pyke CM and Baade PD. The descriptive epidemiology of female breast cancer: an international comparison of screening, incidence, survival and mortality. *Cancer Epidemiology* 2010; 36: 237– 348.
2. Klopffleisch R, Lenze D, Hummel M and Grudlbacher AD. Metastatic canine carcinomas can be identified by a gene expression profile that partly overlaps with human breast cancer profiles. *BMC Cancer* 2010; 10: 618.doi:10.1186/1471-2407-10-618.
3. Lavallo GE, Bertagnoli C, Tavares WLF and Cassali GD. Cox-2 expression in canine mammary carcinomas: correlation angiogenesis and overall survival. *Veterinary Pathology* 2009; 46: 1275– 1280.
4. Laumbacher B, Fellerhoff B, Herzberger B and Wank R. Do dogs harbour risk factors for human breast cancer? *Medical Hypotheses* 2006; 67: 21– 26.
5. Marchetti V, Giorgi M, Fioravanti A, Finotello R, Citi S, Canu B, Orlandi P, Di Desidero T, Danesi R and Bocci G. First-line metronomic chemotherapy in a metastatic model of spontaneous canine tumours: a pilot study. *Investigational New Drugs* 2012; 30: 1725– 1730.
6. Pierini A, Bocci C, Giorgi M, Owen H and Marchetti V. From humans to dogs and back: the transitional lesson of metronomic chemotherapy. *American Journal of Animal and Veterinary Sciences* 2012; 7: 198– 212.

7. Perez Alenza MD, Pen˜a L, del Castillo N and Nieto AI. Factors influencing the incidence and prognosis of canine mammary tumours. *The Journal of Small Animal Practice* 2000; 41: 287– 291.
8. Fitzgibbons PL, Page DL, Weaver D, Thor AD, Allred DC, Clark GM, Ruby SG, O'Malley F, Simpson JF, Connolly JL, Hayes DF, Edge SB, Lichter A and Schnitt SJ. Prognostic factors in breast cancer. College of American Pathologists Consensus Statement 1999. *Archives of Pathology & Laboratory Medicine* 2000; 124: 966– 978.
9. Veronesi B and Oortgiesen M. The TRPV1 receptor: target of toxicants and therapeutics. *Toxicological Sciences* 2006; 89: 1– 3.
10. Chou MZ, Mtui T, Gao Y-D, Kohler M and Middleton RE. Resiniferatoxin binds to the capsaicin receptor (TRPV1) near the extracellular side of the S4 transmembrane domain. *Biochemistry* 2004; 43: 2501– 2511.
11. Richardson JD and Vasko MR. Cellular mechanisms of neurogenic inflammation. *Journal of Pharmacology and Experimental Therapeutics* 2002; 302: 839– 845.
12. Zhang L and Barritt GJ. TRPM8 in prostate cancer cells: a potential diagnostic and prognostic marker with a secretory function? *Endocrine-Related Cancer* 2006; 13: 27– 38.
13. Marincsaˆk R, Toˆth BI, Czifra G, Maˆrton I, Reˆdl P, Tar I, Toˆth L, Kovaˆcs L and Bıroˆ T. Increased expression of TRPV1 in squamous cell carcinoma of the human tongue. *Oral Diseases* 2009; 15: 328– 335.
14. Czifra G, Varga A, Nyeste K, Marincsaˆk R, Toˆth BI, Kovaˆcs I, Kovaˆcs L and Bıroˆ T. Increased expressions of cannabinoid receptor-1 and transient receptor potential vanilloid-1 in human prostate carcinoma. *Journal of Cancer Research and Clinical Oncology* 2009; 135: 507– 514.
15. Vinuesa AG, Sancho R, Garcıa-Limones C, Behrens A, ten Dijke P, Calzado MA and Mun˜oz E. Vanilloid receptor-1 regulates neurogenic inflammation in colon and protects mice from colon cancer. *Cancer Research* 2012; 72: 1705– 1716.
16. Santoni G, Caprodossi S, Farfariello V, Liberati S, Gismondi A, Amantini C. Antioncogenic effects of transient receptor potential vanilloid 1 in the progression of transitional urothelial cancer of human bladder. *ISRN Urology* 2012. doi: 10.5402/2012/458238.
17. Nilius B. TRP channels in disease. *Biochemica and Biophysica Acta* 2007; 1772: 805– 812.
18. Hayes P, Meadows HJ, Gunthorpe MJ, Harries MH, Duckworth DM, Cairns W, Harrison DC, Clarke CE, Ellington K, Prinjha RK, Baarton AJ, Medhurts AD, Smith GD, Topp S, Murdock P, Sanger GJ, Randall AD, Gloger IS and Davis JB. Cloning and functional expression of a human orthologue of rat vanilloid receptor-1. *Pain* 2000; 88: 205– 215.
19. Caterina MJ, Schumaker MA, Tominaga M, Rosen TA, Levine JD and Julius D. The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* 1997; 389: 816– 824.
20. Savidge J, Davis C, Shan K, Calley S, Phillips E, Ranasinghe S, Winter J, Kotsonis P, Rang H and McIntyre P. Cloning and functional characterization of the guinea pig vanilloid receptor

1. Neuropharmacology 2012; 43: 450– 456.
21. Gavva NR, Klionsky L, Qu Y, Shi L, Tamir R, Edenson S, Zhang TJ, Viswanadhan VN, Toeth A, Pearce LV, Vanderah TW, Porreca F, Blumberg PM, Lile J, Sun Y, Wild k, Luis JC, Treanor J. Molecular determinants of vanilloid sensitivity. *Journal of Biological Chemistry* 2004; 279: 20283– 20295.
22. Correll CC, Phelps PT and Anthares JA. Cloning and characterization of mouse TRPV1. *Neuroscience Letters* 2004; 370: 55– 60.
23. Phelps PT, Anthes JC and Correll CC. Cloning and functional characterization of dog transient potential vanilloid receptor-1 (TRPV1). *European Journal of Pharmacology* 2005; 513: 57– 66.
24. Phillips E, Reeve A, Bevan S and McIntyre P. Identification of species-specific determinants of the action of the antagonist capsazepine and the agonist PPAHV on TRPV1. *Journal of Biological Chemistry* 2004; 279: 17165– 17172.
25. McIntyre P, McLatchie LM, Chamebers A, Phillips E, Clarke M, Sadvige J, Toms C, Peacock M, Shah K, Winter J, Weerasakera N, Webb M, Rang MP, Bevan S and James IF. Pharmacological differences

between the human and rat vanilloid receptor 1 (VR1). *British Journal of Pharmacology* 2001; 132: 1084– 1094.

26. Pinho SS, Carvalho S, Cabral J, Reis CA and Gärtner

F. Canine tumors: a spontaneous animal model of human carcinogenesis. *Translational Research* 2012; 159: 165– 172.

27. Shade GR, Keller J, Ives K, Cheng X, Rosol TJ, Keller E and Roberts WW. Histotripsy focal ablation of implanted prostate cancer tumor in an ACE-1 canine cancer model. *Journal of Urology* 2012; 188: 1957– 1964.

28. Maniscalco L, Iussich S, Morello E, Martano M, Biolatti B, Riondato F, Salda LD, Romanucci M, Malatesta D, Bongiovanni L, Tirrito F, Gattino F, Buracco P, DeMaria R. PDGFs and PDGFRs in canine osteosarcoma: new targets for innovative therapeutic strategies in comparative oncology. *Veterinary Journal* 2012; 195: 41– 47.

29. Marconato L, Gelain ME and Comazzi S. The dog as a possible animal model for human non-Hodgkin lymphoma: a review. *Journal of Hematology & Oncology* 2012. doi:10.1002/hon.2017.

30. Peruzzi D, Gavazza A, Mesiti G, Lubas G, Scarselli E, Conforti A, Bendtsen C, Ciliberto G, La Monica N and Aurisicchio L. A vaccine targeting telomerase enhances survival of dogs affected by B-cell lymphoma. *Molecular Therapy* 2010; 18:

1559– 1567.

31. Rivera P and Von Euler H. Molecular biological aspects on canine and human mammary tumors. *Veterinary Pathology* 2011; 48: 132– 146.

32. Lowry OH, Rosebrough NJ, Farr AL and Randall RJ. Protein measurement with the Folin phenol reagent. *Journal of Biological Chemistry* 1951; 193: 265– 275.

33. Abe K and Matsuki N. Measurement of cellular 3- (4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) reduction activity and lactate dehydrogenase release using MTT. *Neuroscience Research* 2000; 38: 325– 329.

34. Bodo´ E, Kova´cs I, Telek A, Varga A, Paus R, Kova´cs L and Biro´ T. Vanilloid receptor-1 (VR1) is widely expressed on various epithelial and mesenchymal cell types of human skin. *Journal of Investigative Dermatology* 2004; 123: 410– 413.
35. Barbero R, Badino P, Cuniberti B, Miolo A, Odore R, Di Marzo V and Re G. Identification of functional TRPV1 vanilloid receptor in MCF-7 cells. *Journal of Veterinary Pharmacology and Therapeutics* 2006; 29(Suppl. 1): 203– 204.
36. Li S, Bode AM, Zhu F, Liu K, Zhang J, Kim MO, Reddy K, Zykova T, Ma WY, Carper AL, Langfald

AK and Dong Z. TRPV1-antagonist AMG9810 promotes mouse skin tumorigenesis through EGFR/Akt signaling. *Carcinogenesis* 2011; 32:

779– 785.

37. Dhennin-Duthille I, Gautier M, Faouzi M, Guilbert A, Brevet M, Vaudry D, Ahidouch A, Sevestre H and Ouadid-Ahidouch H. High expression of transient receptor potential channels in human breast cancer epithelial cells and tissues: correlation with pathological parameters. *Cellular Physiology and Biochemistry* 2011; 28: 813– 822.
38. Ligresti A, Moriello AS, Starowicz K, Matias I, Pisanti S, De Petrocellis L, Laezza C, Portella G, Bifulco M and Di Marzo V. Antitumor activity of plant cannabinoids with emphasis on the effect of cannabidiol on human breast carcinoma. *Journal of Pharmacology and Experimental Therapeutics* 2006; 318: 1375– 1387.
39. Freimuth N, Ramer R and Hinz B. Antitumorigenic effects of cannabinoids beyond apoptosis. *Journal of Pharmacology and Experimental Therapeutics* 2010; 332: 336– 344.
40. Mogg AJ, Mill CE, Folly EA, Beattie RE, Blanco MJ, Beck JP, Broad LM. Altered pharmacology of native rodent spinal cord TRPV1 after phosphorylation. *British Journal of Pharmacology* 2012; 168: 1015– 1029.
41. Hwang MK, Bode AM, Byun S, Song NR, Lee HJ, Lee KW and Dong Z. Cocarcinogenic effect of capsaicin involves activation of EGFR signaling but not TRPV1. *Cancer Research* 2010; 70: 6859– 6869.
42. Jung J, Lee SY, Hwang SW, Cho H, Shin J, Kang YS, Kim S and Oh U. Agonist recognition sites in the cytosolic tails of vanilloid receptor 1. *Journal of Biological Chemistry* 2002; 277: 44448– 44454.
43. Premkumar LS and Sikand P. TRPV1: a target for next generation analgesics. *Current Neuropharmacology* 2008; 6: 151– 163.
44. Tominaga M and Tominaga T. Structure and function of TRPV1. *Pflugers Arch: European Journal of Physiology* 2005; 451: 143– 150.
45. Oyagbemi AA, Saba AB and Azeez OI. Capsaicin: a novel chemopreventive molecule and its underlying molecular mechanisms of action. *Indian Journal of Cancer* 2010; 47: 53– 58.
46. Yang ZH, Wang XH, Wang HP, Hu LQ, Zheng XM and Li SW. Capsaicin mediates cell death in bladder cancer T24 cells through reactive oxygen species production and mitochondrial depolarization. *Urology* 2010; 75: 735– 741.
47. Gil YG and Kang MK. Capsaicin induces apoptosis and terminal differentiation in human glioma A172 cells. *Life Sciences* 2008; 82: 997– 1003.